

1

WhisperJECT™ Autoinjector*

Intended for use with FDA-approved drug products which are presented in a 1.0 mL BD®, pre-filled, glass syringe with a fixed needle of 27G to 29G

2

1.0 mL glass pre-filled syringe

Additional supplies you need:

- A clean damp cloth
- A sharps container
- A place to record your injection

3

Alcohol
Wipe

4

Cotton
Ball

Please see Important Safety Information on the following page. For injection instructions, please see the full Prescribing Information, including Patient Information Leaflet and Instructions for Use. Please see the Instructions for Use provided with your WhisperJECT™ Autoinjector device.

Call **1.844.695.2667**
for 24/7 nurse support
regarding your injection

Mylan's Glatiramer Acetate Injection is available by prescription only.

*WhisperJECT™ Autoinjector is available by Rx only.

Mylan's Glatiramer Acetate Injection logo is a trademark of Mylan Pharmaceuticals Inc.

WhisperJECT™ is a trademark of Mylan Pharmaceuticals Inc.

The Mylan logo is a registered trademark of Mylan Inc.

BD® is a registered trademark of Becton, Dickinson and Company.

©2017 Mylan Pharmaceuticals Inc. All rights reserved. GA-2017-0017

INDICATION

GLATIRAMER ACETATE INJECTION is a prescription medicine used for the treatment of people with relapsing forms of multiple sclerosis (MS).

IMPORTANT SAFETY INFORMATION

Do not take GLATIRAMER ACETATE INJECTION if you are allergic to glatiramer acetate or mannitol.

Some patients report a short-term reaction right after injecting glatiramer acetate. This reaction can involve flushing (feeling of warmth and/or redness), chest tightness or pain with heart palpitations, anxiety, and trouble breathing. These symptoms generally appear within minutes of an injection, last about 15 minutes, and do not require specific treatment. There have been reports of patients with similar symptoms who received emergency medical care. **If symptoms become severe, call the emergency phone numbers in your area.**

Call your doctor right away if you develop hives, skin rash with irritation, dizziness, sweating, chest pain, trouble breathing, or severe pain at the injection site. If any of the above occurs, do not give yourself any more injections until your doctor tells you to begin again.

Chest pain may occur either as part of the immediate post-injection reaction or on its own. This pain should only last a few minutes. You may experience more than one such episode, usually beginning at least one month after starting treatment. Tell your doctor if you experience chest pain that lasts for a long time or feels very intense.

A permanent indentation under the skin (lipoatrophy and, rarely, death of your skin tissue also referred to as necrosis) at the injection site may occur due to local destruction of fat tissue. Be sure to follow proper injection technique and inform your doctor of any skin changes.

The most common side effects in studies of GLATIRAMER ACETATE INJECTION are redness, pain, swelling, itching, or a lump at the site of injection, flushing, rash, shortness of breath, and chest pain. These are not all of the possible side effects of GLATIRAMER ACETATE INJECTION. For a complete list, ask your doctor or pharmacist. Tell your doctor about any side effects that you have while taking GLATIRAMER ACETATE INJECTION.

You are encouraged to report negative side effects of prescription drugs to the FDA.

Visit www.fda.gov/medwatch or call **1-800-FDA-1088**.

Please see full Prescribing Information, Patient Information Leaflet and Instructions for Use for [GLATIRAMER ACETATE INJECTION 20 mg/mL](#) or [GLATIRAMER ACETATE INJECTION 40 mg/mL](#).