

1

WhisperJECT[®]
AUTOINJECTOR

Intended for use with FDA-approved drug products which are presented in a 1.0 mL BD[®], pre-filled, glass syringe with a fixed needle of 27G to 29G

2

1.0 mL glass pre-filled syringe

Additional supplies you need:

- A clean damp cloth
- A sharps container
- A place to record your injection

3

Alcohol
Wipe

4

Cotton
Ball

Please see Important Safety Information on the following page. For injection instructions, please see the full Prescribing Information, including Patient Information Leaflet and Instructions for Use. Please see the Instructions for Use provided with your *WhisperJECT*[®] Autoinjector device.

Call 1.844.695.2667
for 24/7 nurse support
for help with your injection

Viatrix' Glatiramer Acetate Injection is available by prescription only.

**WhisperJECT*[®] Autoinjector is available by Rx only.

WhisperJECT and the Glatiramer Acetate Injection Logo are registered trademarks of Mylan Pharmaceuticals Inc., a Viatrix Company.

VIATRIS and the Viatrix Logo are trademarks of Mylan Inc., a Viatrix Company.

BD[®] is a registered trademark of Becton, Dickinson and Company.

© 2020 Viatrix Inc. All Rights Reserved. GA-2019-0115 V4

USE

GLATIRAMER ACETATE INJECTION is a prescription medicine used for the treatment of people with relapsing forms of multiple sclerosis (MS), to include clinically isolated syndrome, relapsing-remitting disease, and active secondary progressive disease, in adults.

IMPORTANT SAFETY INFORMATION

Do not take GLATIRAMER ACETATE INJECTION if you are allergic to glatiramer acetate or mannitol.

Serious side effects may happen right after or within minutes after you inject glatiramer acetate injection at any time during your course of treatment. Call your doctor right away if you have any of these immediate post-injection reaction symptoms including: redness to your cheeks or other parts of the body (flushing); chest pain; fast heart beat; anxiety; breathing problems or tightness in your throat; or swelling, rash, hives, or itching. If you have symptoms of an immediate post-injection reaction, do not give yourself more injections until a doctor tells you to.

Chest pain may occur either as part of the immediate post-injection reaction or on its own. This pain usually only lasts a few minutes or can begin around 1 month after you start using glatiramer acetate injection. Call your doctor right away if you experience chest pain.

A permanent indentation under the skin (lipoatrophy and, rarely, death of your skin tissue also referred to as necrosis) at the injection site may occur due to local destruction of fat tissue. In order to reduce your chance of developing these problems, be sure to follow your doctor's instruction on how to use glatiramer acetate injection and be sure to choose a different injection site each time you use glatiramer acetate injection.

Liver problems, including liver failure, can occur with GLATIRAMER ACETATE INJECTION. Call your healthcare provider right away if you have symptoms, such as nausea, loss of appetite, tiredness, dark colored urine and pale stools, yellowing of your skin or the white part of your eye, bleeding more easily than normal, confusion, or sleepiness.

The most common side effects in studies of GLATIRAMER ACETATE INJECTION are redness, pain, swelling, itching, or a lump at the site of injection, rash, shortness of breath, and flushing. These are not all the possible side effects of GLATIRAMER ACETATE INJECTION. For a complete list, ask your doctor or pharmacist. Tell your doctor about any side effects that you have while taking GLATIRAMER ACETATE INJECTION.

You are encouraged to report negative side effects of prescription drugs to the FDA.

Visit www.fda.gov/medwatch or call **1-800-FDA-1088**.

Please see full Prescribing Information, Patient Information Leaflet and Instructions for Use for [GLATIRAMER ACETATE INJECTION 20 mg/mL](#) or [GLATIRAMER ACETATE INJECTION 40 mg/mL](#).