



While refrigeration is preferred, when you are not able to refrigerate GLATIRAMER ACETATE INJECTION, you may store it for up to 1 month at room temperature between 15° to 30° C (59° to 86° F).

Treating relapsing forms of **Multiple Sclerosis (MS)**, to include clinically isolated syndrome, relapsing-remitting disease, and active secondary progressive disease, in adults

AVAILABLE IN 2 DOSAGE STRENGTHS:

20 mg/mL

for **once-daily** injection

40 mg/mL

for **3-times-a-week** injection

USE

GLATIRAMER ACETATE INJECTION is a prescription medicine that is used to treat relapsing forms of multiple sclerosis (MS), to include clinically isolated syndrome, relapsing-remitting disease, and active secondary progressive disease, in adults. It is not known if GLATIRAMER ACETATE INJECTION is safe and effective in children under 18 years of age.

IMPORTANT SAFETY INFORMATION

Do not take GLATIRAMER ACETATE INJECTION if you are allergic to glatiramer acetate or mannitol.

[Click here](#) for additional Important Safety Information on page 25 and 26 of this brochure.



Viatrix'

COMMITMENT TO YOU

You can depend on the VIATRIS ADVOCATE® patient support program to help you access therapy as soon as possible and provide you with an array of support services to assist you in achieving your goals.

In addition, VIATRIS™ offers you our *WhisperJECT*® Autoinjector* device, intended for use with FDA-approved drug products with non-viscous (aqueous) liquid formulations, which are presented in a 1.0 mL BD®, pre-filled, glass syringe with staked needle, containing a fixed needle of 27G to 29G. The *WhisperJECT* Autoinjector is a reusable injection device for the subcutaneous injection of certain FDA-approved drugs, such as Viatrix' Glatiramer Acetate Injection. *WhisperJECT* is a device with features that have been designed by patients, for patients.

We are one of the world's leading generic and specialty pharmaceuticals companies. Our dedication to providing access to important medicine for the world's 7 billion people continues to grow.

IMPORTANT SAFETY INFORMATION (CONTINUED)

Serious side effects may happen right after or within minutes after you inject GLATIRAMER ACETATE INJECTION at any time during your course of treatment. Call your healthcare provider right away if you have any of these immediate post-injection reaction symptoms including redness to your cheeks or other parts of the body (flushing); chest pain; fast heartbeat; anxiety; breathing problems or tightness in your throat; or swelling, rash, hives, or itching.

**WhisperJECT* Autoinjector is available by Rx only.







VIATRIS
ADVOCATE

**Learn more about VIATRIS ADVOCATE
on page 8 of this brochure.**

Before You Begin

You should receive your first dose of Viatrix' Glatiramer Acetate Injection under the direction of a doctor or nurse, whether administered with the pre-filled syringe or autoinjector. This might be at your doctor's office or a nurse from VIATRIS ADVOCATE can teach you how to give your glatiramer acetate injection in person or via video conference or telephone.

Carefully read the Patient Information and Instructions for Use sections in the Full Prescribing Information for Viatrix' Glatiramer Acetate Injection, provided with this brochure.

If you use the *Whisper*JECT Autoinjector device, please read the Instructions for Use inside the carrying case.

IMPORTANT SAFETY INFORMATION (CONTINUED)

If you have symptoms of an immediate post-injection reaction, do not give yourself more injections until a healthcare provider tells you to.

Chest pain may occur either as part of the immediate post-injection reaction or by itself. This type of chest pain usually lasts a few minutes and can begin around 1 month after you start using GLATIRAMER ACETATE INJECTION. Call your healthcare provider right away if you experience chest pain.

Damage to the fatty tissue just under your skin's surface (lipoatrophy) and, rarely, death of your skin tissue (necrosis) can happen when you use glatiramer acetate injection. Damage to the fatty tissue under your skin can cause a "dent" at the injection site that may not go away. You can reduce your chance of developing these problems by:

- following your healthcare provider's instructions for how to use glatiramer acetate injection
- choosing a different injection area each time you use glatiramer acetate injection

Helpful Resources

You may also find helpful information in the Guide for Self-Injection With *WhisperJECT Autoinjector (Rx only)* and the *Injection Help* section of the website, with more details about self-injection.


You can download a *Preparation Mat* for your injection, with designated spaces for your supplies, and a *Rotation Plan*, where you can record the area, site, day and depth setting (of the *WhisperJECT Autoinjector* device) of each injection, at [GlatiramerAcetate.com](https://www.glatirameracetate.com).



The information in this brochure does not take the place of talking with your healthcare provider about your medical condition or treatment.

IMPORTANT SAFETY INFORMATION (CONTINUED)

Liver problems, including liver failure, can occur with GLATIRAMER ACETATE INJECTION. Call your healthcare provider right away if you have symptoms, such as nausea, loss of appetite, tiredness, dark colored urine and pale stools, yellowing of your skin or the white part of your eye, bleeding more easily than normal, confusion, or sleepiness.



Identify symptoms, including any that vary by time of day or by position/activity.



Location	Day	Onset/Setting
Front of Upper Right Arm		
Back of Upper Right Arm		
Front of Upper Left Arm		
Back of Upper Left Arm		
Front of Upper Right Leg		
Back of Upper Right Leg		
Front of Upper Left Leg		
Back of Upper Left Leg		
Front of Right Thigh		
Back of Right Thigh		
Front of Left Thigh		
Back of Left Thigh		
Abdomen		

Remember: There is a yellow band on the upper right of the "Function" area.

Download your free
Preparation Mat and Rotation Plan at
GlatiramerAcetate.com



Here when you need us!

(1.844.695.2667)



Not an actual nurse.

VIATRIS ADVOCATE is your support team

- a program to help you get started on and stay committed to your doctor's treatment plan for Viatris' Glatiramer Acetate Injection. Whether you have questions for our MS-experienced nurses, or need assistance from our health benefits experts with your insurance coverage or copay assistance, VIATRIS ADVOCATE is there for you. Our nurses are trained and knowledgeable about your therapy. These specialists can visit you in person, speak with you virtually or by telephone to help you develop a plan for giving your own injections, based on your doctor's direction.

IMPORTANT SAFETY INFORMATION (CONTINUED)

Some glatiramer acetate products can be used with an optional compatible autoinjector. Compatible autoinjectors are supplied separately if available, but the availability of compatible autoinjectors may change with time.


Your VIATRIS ADVOCATE Welcome Pack

Once enrolled in the VIATRIS ADVOCATE program, you will not need to take any further steps to participate in your patient support program. And, you will receive a Welcome Pack consisting of:

- 1 Neoprene Sleeve**
that showcases inspiring art created by Shana Stern, an artist living with a relapsing form of MS
- 2 Patient Brochure**
that highlights Viatris' Glatiramer Acetate Injection, as well as the resources and services available from VIATRIS ADVOCATE
- 3 Guide for Self-Injection**
with useful information about the *WhisperJECT* Autoinjector device and helpful tips for self-injecting
- 4 Needle Clipper**
- 5 Hot/Cold Gel Pack**
- 6 VIATRIS ADVOCATE Magnet**
- 7 Full Prescribing Information**
for Viatris' Glatiramer Acetate Injection, including the Patient Information Leaflet and Instructions for Use

IMPORTANT SAFETY INFORMATION (CONTINUED)

Check with your healthcare provider when you fill or refill your medicine to make sure the autoinjector you have is meant to be used with your glatiramer acetate product. If you use the wrong autoinjector, you might not get the correct dose of your medicine.



Not an actual nurse.

Nurse Support

In addition to the care and direction provided by your doctor, VIATRIS ADVOCATE nurses are experienced in caring for people like you who have relapsing forms of multiple sclerosis. They are available to you by phone at **1.844.695.2667** from Monday through Friday 8 a.m. to 8 p.m. central time, to answer your questions about:

- Your condition
- Viatrix' Glatiramer Acetate Injection
- Proper injection technique

Nurses provide ongoing support, education and encouragement to you, your family and your caregivers.

IMPORTANT SAFETY INFORMATION (CONTINUED)

The most common side effects in studies of GLATIRAMER ACETATE INJECTION are skin problems at your injection site, including redness, pain, swelling, itching, or a lump at the site of injection, rash, shortness of breath, flushing (vasodilation), and chest pain.

Experienced nurses work with you one on one to:

- Go through the injection process step by step to help you learn proper injection techniques
- Help you adjust your routine based on lifestyle changes, such as changes in weight
- Provide you with tips regarding preparation, administration, storage and disposal of your medicine
- Offer you tools for tracking self-injections
- Connect you to supportive MS resources and encourage you to participate in the MS community
- Assist you in connecting to a support network

Injection Training

You can call VIATRIS ADVOCATE to schedule an in-person, virtual, or telephonic training session with one of our MS-experienced nurses, who can show you how to use the pre-filled glass syringe by itself or with the *Whisper*JECT Autoinjector and guide you through each step of your injection.

By staying close with VIATRIS ADVOCATE, you'll find support and services that you need. On the following pages are some of the features and benefits.

IMPORTANT SAFETY INFORMATION (CONTINUED)

These are not all the possible side effects of GLATIRAMER ACETATE INJECTION. For a complete list, ask your healthcare provider or pharmacist. Tell your healthcare provider about any side effects that you have while taking GLATIRAMER ACETATE INJECTION.

Patient Access Specialist

Your healthcare professional is your go-to resource for your treatment plan, and your patient access specialist is your go-to resource for navigating through the requirements of your insurance plan, whether it's private or government sponsored. Your patient access specialist can help you:

- Coordinate and optimize your insurance benefits
- Fill out and file the right forms at the right time
- Sign you up for the VIATRIS ADVOCATE Copay Program, if you're eligible
- Find other potential sources of financial assistance


MS Resources

Following enrollment, VIATRIS ADVOCATE will provide you with information on topics most relevant to you, including brochures, websites and connections to MS community organizations and events. The goal of VIATRIS ADVOCATE is to help MS patients receive the support they need.

IMPORTANT SAFETY INFORMATION (CONTINUED)

Do not take GLATIRAMER ACETATE INJECTION if you are allergic to glatiramer acetate or mannitol.

Serious side effects may happen right after or within minutes after you inject GLATIRAMER ACETATE INJECTION at any time during your course of treatment. Call your healthcare provider right away if you have any of these immediate post-injection reaction symptoms including redness to your cheeks or other parts of the body (flushing); chest pain; fast heartbeat; anxiety; breathing problems or tightness in your throat; or swelling, rash, hives, or itching.



Get started by calling **VIATRIS ADVOCATE**
at **1.844.695.2667** today.



Not an actual patient.

VIATRIS ADVOCATE Copay Program

For eligible, commercially insured patients

Your copay for Viatrix' Glatiramer Acetate Injection could be as low as \$0 per month.*

- No financial qualifications
- If your insurance recently changed, contact VIATRIS ADVOCATE to see if you are eligible for our copay program

* See eligibility criteria and full terms and conditions on the next page.

For the full terms and conditions, please see

www.glatirameracetate.com/en/savings-support

IMPORTANT SAFETY INFORMATION (CONTINUED)

If you have symptoms of an immediate post-injection reaction, do not give yourself more injections until a healthcare provider tells you to.



Eligibility Criteria Terms and Conditions

- With this VIATRIS ADVOCATE copay assistance program, patients may pay as little as \$0 for each monthly fill of VIATRIS' Glatiramer Acetate Injection 20 mg/mL and/or 40 mg/mL, while this program remains in effect. This copay assistance program may be used to reduce the amount of an eligible patient's out-of-pocket costs for VIATRIS' Glatiramer Acetate Injection 20 mg/mL and/or 40 mg/mL up to the full amount of the patient's out-of-pocket cost per prescription, which may be up to a 90-day supply, subject to a maximum aggregate amount of \$9,000 per 12-month period while this copay assistance program remains in effect (such aggregate amount includes dispenses of both VIATRIS' Glatiramer Acetate Injection 20 mg/mL and 40 mg/mL). No other purchase is necessary. Valid prescription is required. VIATRIS reserves the right to amend or end this copay assistance program at any time without notice.
- **Eligibility Requirements: This copay assistance can be redeemed only by patients or patient guardians who are 18 years of age or older and who are residents of the U.S. or Puerto Rico. Patients must have commercial prescription drug insurance. This copay assistance program is not valid for uninsured patients (but may be used by commercially insured patients without coverage for VIATRIS' Glatiramer Acetate Injection 20 mg/mL and/or 40 mg/mL) and patients who are covered in whole or in part by any state or federally funded healthcare program, including, but not limited to, any state pharmaceutical assistance program, Medicare (Part D or otherwise), Medicaid, Medigap, VA or DOD, or TRICARE (regardless of whether VIATRIS' Glatiramer Acetate Injection 20 mg/mL and/or 40 mg/mL is covered by such government program); not valid if the patient is Medicare eligible and enrolled in an employer-sponsored health plan or prescription benefit program for retirees; and not valid if the patient's insurance plan is paying the entire cost of this prescription. This copay assistance program is void outside the U.S. or Puerto Rico or in any state or jurisdiction where prohibited by law, taxed or restricted. Absent a change in Massachusetts law, this copay assistance program will no longer be valid for**
 - **Massachusetts residents as of January 1, 2026. This copay assistance program is not valid for residents of California.**
 - This copay assistance program is not health insurance. The copay assistance program is not transferable, and the amount of the savings cannot exceed the patient's out-of-pocket costs. Cannot be combined with any other rebate/coupon, cash discount card, free trial, or similar offer for the specified prescription. This copay assistance is not redeemable for cash. This copay assistance is not valid for product dispensed by a 340B covered entity that purchased the product at discounted pricing under the 340B drug pricing program. This copay assistance is not valid if the patient's commercial health insurance plan or pharmacy benefit manager uses a copay adjustment program (often termed "maximizer" or "accumulator" program) that restricts any form of copay assistance from being counted toward the patient's cost-sharing limits.
 - **NOTICE:** Data related to your use of this copay assistance program may be collected, analyzed and shared with VIATRIS for market research and other purposes related to assessing its copay assistance programs. Data shared with VIATRIS will be aggregated and de-identified, meaning it will be combined with data related to other copay assistance program redemptions and will not identify you.
 - Use of this copay assistance program must be consistent with the terms of any drug benefit provided by a commercial health insurer, health plan or private third-party payer. Patients must have not submitted and will not submit a claim for reimbursement under any federal, state or other governmental programs for this prescription. Patients are responsible for reporting the receipt of copay assistance to any commercial insurer, health plan, or third-party payer who pays for or reimburses any part of the prescription filled, as may be required. Patients should not use this copay assistance program if their health plan prohibits use of manufacturer copay assistance programs. Patients should withdraw from this copay assistance program should they begin to receive prescription benefits from any government funded program.

IMPORTANT SAFETY INFORMATION (CONTINUED)

Chest pain may occur either as part of the immediate post-injection reaction or by itself. This type of chest pain usually lasts a few minutes and can begin around 1 month after you start using GLATIRAMER ACETATE INJECTION. Call your healthcare provider right away if you experience chest pain.



Not an actual patient.

Storing Viatrix' Glatiramer Acetate Injection

When you receive your supply of Viatrix' Glatiramer Acetate Injection pre-filled syringes, you should mark the date you receive them on the carton and then refrigerate them at 36°-46° F. Ensure they are kept out of the reach of children.

Viatrix' Glatiramer Acetate Injection should never be frozen or exposed to intense light. If a glatiramer acetate injection syringe freezes, it should be discarded in a sharps disposal container.

While refrigeration is preferred, you may keep Viatrix' Glatiramer Acetate Injection at room temperature (59°-86° F) for up to 1 month when you do not have access to refrigeration.

IMPORTANT SAFETY INFORMATION (CONTINUED)


Damage to the fatty tissue just under your skin's surface (lipoatrophy) and, rarely, death of your skin tissue (necrosis) can happen when you use glatiramer acetate injection. Damage to the fatty tissue under your skin can cause a "dent" at the injection site that may not go away. You can reduce your chance of developing these problems by:

- following your healthcare provider's instructions for how to use glatiramer acetate injection
- choosing a different injection area each time you use glatiramer acetate injection

Features of the *Whisper*JECT

Viatrix has developed the *Whisper*JECT Autoinjector,* which is intended for use with FDA-approved drug products with non-viscous (aqueous) liquid formulations, which are presented in a 1.0 mL BD, pre-filled glass syringe with staked needle, containing a fixed needle of 27G to 29G. The *Whisper*JECT Autoinjector is a reusable injection device for the subcutaneous injection of certain FDA-approved drugs such as Viatrix' Glatiramer Acetate Injection.

The *Whisper*JECT Autoinjector device has features that have been designed by patients, for patients:



If you use the *Whisper*JECT Autoinjector device, please read the Instructions for Use inside the carrying case.

**Whisper*JECT Autoinjector is available by Rx only.

Check with your healthcare provider when you fill or refill your medicine to make sure the autoinjector you have is meant to be used with your glatiramer acetate product. If you use the wrong autoinjector, you might not get the correct dose of your medicine.



Not an actual nurse.

Caring for Your *WhisperJECT* Autoinjector

Cleaning your *WhisperJECT* Autoinjector device is easy. Wipe the accessible areas with a clean, damp cloth after each use. Do not immerse in water and do not clean with solvents, detergent or alcohol.

Your device needs to be replaced after 2 years.

Contact VIATRIS ADVOCATE at 1.844.695.2667 for replacement.

IMPORTANT SAFETY INFORMATION (CONTINUED)

Liver problems, including liver failure, can occur with GLATIRAMER ACETATE INJECTION. Call your doctor right away if you have any of these immediate post-injection reaction symptoms Call your healthcare provider right away if you have symptoms, such as nausea, loss of appetite, tiredness, dark colored urine and pale stools, yellowing of your skin or the white part of your eye, bleeding more easily than normal, confusion, or sleepiness.

Traveling with Viatrix' Glatiramer Acetate Injection and the *WhisperJECT* Autoinjector

You can travel with Viatrix' Glatiramer Acetate Injection.

Pre-filled syringes may be kept at room temperature (59°-86° F) for up to 1 month during travel or when refrigeration may not be available.

- Make sure to bring enough supplies with you. In case of emergencies or a change in plans, pack more syringes than you need. Be sure to pack enough cotton balls and alcohol wipes for the number of days you'll be away
- You might also want to use a tight-locking plastic storage bag for all of your supplies
- If you're traveling by plane, keep Viatrix' Glatiramer Acetate Injection pre-filled syringes and the *WhisperJECT* Autoinjector device in its travel case in your carry-on luggage, so they're easily accessible
- To avoid unnecessary delays, bring your current prescription label with you; your name on the prescription label should match your name as it appears on your ID



For current regulations, contact the Transportation Security Administration, or visit their website at [tsa.gov](https://www.tsa.gov).

IMPORTANT SAFETY INFORMATION (CONTINUED)

Some glatiramer acetate products can be used with an optional compatible autoinjector. Compatible autoinjectors are supplied separately if available, but the availability of compatible autoinjectors may change with time.



Not an actual patient.

IMPORTANT SAFETY INFORMATION (CONTINUED)

Check with your healthcare provider when you fill or refill your medicine to make sure the autoinjector you have is meant to be used with your glatiramer acetate product. If you use the wrong autoinjector, you might not get the correct dose of your medicine.

The most common side effects in studies of GLATIRAMER ACETATE INJECTION are skin problems at your injection site, including redness, pain, swelling, itching, or a lump at the site of injection, rash, shortness of breath, flushing (vasodilation), and chest pain.

Injection Tips

For detailed instructions on how to use Viatris' Glatiramer Acetate Injection, please see the Instructions for Use in the Full Prescribing Information, the *Guide for Self-Injection* as well as the *Injection Help* section of [GlatiramerAcetate.com](https://www.glatirameracetate.com).

Here are some helpful tips to remember each time you self-inject:

- You should remove your pre-filled glass syringe from the refrigerator about 20 minutes before you self-inject
- You can lay out your supplies for injection on the *Preparation Mat*, which you can download from [GlatiramerAcetate.com](https://www.glatirameracetate.com)
- Choose an area of your home to inject that is well lighted and has a clean, flat surface
- Always wash your hands thoroughly with soap and water before injecting. Be careful not to touch your face or hair after washing your hands
- Viatris' Glatiramer Acetate Injection is given by injection under your skin (subcutaneously). Do not inject Viatris' Glatiramer Acetate Injection in your veins (intravenously)
- If using an optional autoinjector for administration, ensure the device is compatible for use with Viatris' Glatiramer Acetate Injection


IMPORTANT SAFETY INFORMATION (CONTINUED)

These are not all the possible side effects of GLATIRAMER ACETATE INJECTION. For a complete list, ask your healthcare provider or pharmacist. Tell your healthcare provider about any side effects that you have while taking GLATIRAMER ACETATE INJECTION.



Click here for the *Guide for Self-injection* and [GlatiramerAcetate.com](https://www.GlatiramerAcetate.com) for detailed information about your injections.

Inject Glatiramer Acetate into the fatty layer under your skin (subcutaneously).



Remember: There are multiple injection sites within each of the 7 injection areas.

Location	Day	Depth Setting
Back of Upper Right Arm (fleshy area)		
Back of Upper Left Arm (fleshy area)		
Back of Upper Right Hip (below waist)		
Back of Upper Left Hip (below waist)		
Front of Right Thigh (2 inches above knee and 2 inches below groin)		
Front of Left Thigh (2 inches above knee and 2 inches below groin)		
Abdomen (Stomach area; avoid 2 inches around the belly button)		

Injection Site Rotation Plan

It's important to keep up with your injections of Viatrix' Glatiramer Acetate Injection, whether you're at home or on the go. Viatrix has tools to help you, including an injection *Rotation Plan* you can download from our website, GlatiramerAcetate.com.

If prescribed the *Whisper*JECT Autoinjector device, you can manually record the area, day, site and depth setting for each injection.

IMPORTANT SAFETY INFORMATION (CONTINUED)

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

USE

GLATIRAMER ACETATE INJECTION is a prescription medicine that is used to treat relapsing forms of multiple sclerosis (MS), to include clinically isolated syndrome, relapsing-remitting disease, and active secondary progressive disease, in adults. It is not known if GLATIRAMER ACETATE INJECTION is safe and effective in children under 18 years of age.

IMPORTANT SAFETY INFORMATION

Do not take GLATIRAMER ACETATE INJECTION if you are allergic to glatiramer acetate or mannitol.

Serious side effects may happen right after or within minutes after you inject GLATIRAMER ACETATE INJECTION at any time during your course of treatment. Call your healthcare provider right away if you have any of these immediate post-injection reaction symptoms including redness to your cheeks or other parts of the body (flushing); chest pain; fast heartbeat; anxiety; breathing problems or tightness in your throat; or swelling, rash, hives, or itching. If you have symptoms of an immediate post-injection reaction, do not give yourself more injections until a healthcare provider tells you to.

Chest pain may occur either as part of the immediate post-injection reaction or by itself. This type of chest pain usually lasts a few minutes and can begin around 1 month after you start using GLATIRAMER ACETATE INJECTION. Call your healthcare provider right away if you experience chest pain.

Damage to the fatty tissue just under your skin's surface (lipoatrophy) and, rarely, death of your skin tissue (necrosis) can happen when you use glatiramer acetate injection. Damage to the fatty tissue under your skin can cause a "dent" at the injection site that may not go away. You can reduce your chance of developing these problems by:

- following your healthcare provider's instructions for how to use glatiramer acetate injection

IMPORTANT SAFETY INFORMATION (CONTINUED)

- choosing a different injection area each time you use glatiramer acetate injection

Liver problems, including liver failure, can occur with GLATIRAMER ACETATE INJECTION. Call your healthcare provider right away if you have symptoms, such as nausea, loss of appetite, tiredness, dark colored urine and pale stools, yellowing of your skin or the white part of your eye, bleeding more easily than normal, confusion, or sleepiness.

Some glatiramer acetate products can be used with an optional compatible autoinjector. Compatible autoinjectors are supplied separately if available, but the availability of compatible autoinjectors may change with time. Check with your healthcare provider when you fill or refill your medicine to make sure the autoinjector you have is meant to be used with your glatiramer acetate product. If you use the wrong autoinjector, you might not get the correct dose of your medicine.

The most common side effects in studies of GLATIRAMER ACETATE INJECTION are skin problems at your injection site, including redness, pain, swelling, itching, or a lump at the site of injection, rash, shortness of breath, flushing (vasodilation), and chest pain. These are not all the possible side effects of GLATIRAMER ACETATE INJECTION. For a complete list, ask your healthcare provider or pharmacist. Tell your healthcare provider about any side effects that you have while taking GLATIRAMER ACETATE INJECTION.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Please click below for Full Prescribing Information, including Patient Information Leaflet and Instructions for Use, for GLATIRAMER ACETATE INJECTION 20 mg/mL or GLATIRAMER ACETATE INJECTION 40 mg/mL

WhisperJECT and the Glatiramer Acetate Injection Logo are registered trademarks of Mylan Pharmaceuticals Inc., a Viatrix Company.

VIATRIS ADVOCATE and the Viatrix Advocate Logo are registered trademarks of Mylan Inc., a Viatrix Company.

VIATRIS and the Viatrix Logo are trademarks of Mylan Inc., a Viatrix Company.

BD® is a registered trademark of Becton, Dickinson and Company.

© 2024 Viatrix Inc. All Rights Reserved. GA-2021-0096 V6

For more information, please visit [Viatrix.com](https://www.viatrix.com).



VIATRIS™