

PATIENT INFORMATION (Please print)

Name (First, MI, Last, Suffix): _____
 Date of Birth: _____ Gender: M F
 Home Address: _____
 City: _____ State: _____ Zip: _____
 Home Phone: _____ Cell Phone: _____
 Email Address: _____
 Allergies: _____ Other Medications: _____

Additional opt-in services available:

- I would like to receive text message injection reminders and I have read and agree to the section below titled "Contact by telephone or text."
- I would like to receive periodic phone calls from a VIATRIS ADVOCATE® nurse to provide additional therapy support and help answer any therapy-related questions.
- I would like to receive marketing emails per the attached **Patient Marketing Consent Section C.**

PRESCRIBER INFORMATION

Physician: _____
 NP/PA (if prescriber): _____
 Facility Name: _____
 Address: _____
 City: _____ State: _____ Zip: _____
 Phone: _____ Fax: _____
 Office Contact: _____ Email: _____

PRESCRIBER SIGNATURE REQUIRED FOR PRESCRIPTION ORDERS:

Statement of Medical Necessity: Primary Diagnosis ICD-10 CM G35 Treatment of Relapsing Forms of Multiple Sclerosis. I authorize VIATRIS ADVOCATE to provide any information on this form to the insurer of the named patient and to forward the above prescription, by fax or by other mode of delivery, to the pharmacy.

Prescriber's Signature: _____

 (Dispense as Written)

 (Brand Exchange Permissible)

 (NPI#)

 (Date)

INSURANCE INFORMATION

(Attach a copy of patient's insurance card, front & back)

Insurance information not necessary if Prescriber is ordering device and/or injection training only.
 Primary Insurance Name: _____
 Medicare: A B D (attach a copy of red, white and blue Medicare card)
 Beneficiary/Cardholder Name: _____
 Primary Insurance ID#: _____ Group#: _____
 Primary Insurance Phone#: _____
 Does the patient have a pharmacy benefit card? Yes No

By signing below, I have read and agree to the attached **Patient Authorization Section A. (Signature required if Prescriber is ordering VIATRIS™ Glatiramer Acetate Injection)**

X

 Patient/Legal Guardian Signature

 Date (MM/DD/YYYY)

PRESCRIPTION INFORMATION

(Check the box for prescriptions/orders required: Product, Device and/or Injection Training)

- Viatris' Glatiramer Acetate Injection 20 mg/mL pre-filled syringes
 - Inject 20 mg SQ once a day
 - Dispense: 1 box of 30 syringes (30-day supply)
 - May dispense up to a 90-day supply
 - Refills: x 1 year
- Viatris' Glatiramer Acetate Injection 40 mg/mL pre-filled syringes
 - Inject 40 mg SQ 3 times a week
 - Dispense: 1 box of 12 syringes (28-day supply)
 - May dispense up to a 84-day supply
 - Refills: x 1 year

- Whisper*JECT® Autoinjector* device (free of charge)†
 - Use as directed
 - Dispense: 1 device with instructions for use and travel case
 - Refills: None

- VIATRIS ADVOCATE to coordinate injection training (injection training to be performed by a Registered Nurse) Please indicate strength for RN (check the box below).
 - Viatris' Glatiramer Acetate Injection 20 mg/mL pre-filled syringes
 - Viatris' Glatiramer Acetate Injection 40 mg/mL pre-filled syringes

Signature stamps not acceptable.

Please attach all prescriptions on Official State Prescription form if mandated by individual state laws.

Please see the next page for Indication and Important Safety Information, and please see accompanying Full Prescribing Information, including Boxed WARNING, Patient Information Leaflet and Instructions for Use for **GLATIRAMER ACETATE INJECTION 20 mg/mL** or **GLATIRAMER ACETATE INJECTION 40 mg/mL**.

**Whisper*JECT® Autoinjector is available by prescription only. †No party may submit an insurance claim or other claim for payment to any third-party payor (private or government) for this device or training.



Patient Enrollment Form

Phone: 844.695.2667 • Fax: 844.292.8395



A. PATIENT AUTHORIZATION: I authorize my healthcare providers and health insurers to disclose to Viatris Inc. d/b/a VIATRIS ADVOCATE, its affiliates, its program administrator, and their respective agents and service providers (collectively, “VIATRIS ADVOCATE”) my protected health information (“PHI”), including information about my insurance, prescriptions, medical condition and health, so that VIATRIS ADVOCATE may use the information to assist me with benefits support in connection with my treatment with Viatris products, communicate with me regarding such treatment and support, to conduct market research and inform me of treatment alternatives. I understand that once disclosed pursuant to this authorization, my PHI may no longer be protected by federal law and could be re-disclosed to others, but I also understand that VIATRIS ADVOCATE intends to safeguard my PHI and to use and disclose it only for the purposes described herein. I understand that I do not need to sign this authorization in order to receive healthcare treatment or insurance benefits, and that I may cancel the authorization at any time by sending a written notice of cancellation by mail to: VIATRIS ADVOCATE Opt-out Administrator, 1000 Mylan Blvd., Canonsburg, PA 15317, or by fax to 1.844.292.8395. If I do not cancel it, the authorization will remain in effect for five years from the date of my signature on the previous page. I understand that I have a right to receive a copy of this authorization when it is signed.

B. CONTACT BY TELEPHONE OR TEXT:

I agree to be contacted by VIATRIS ADVOCATE by mail, email, telephone calls, and text messages at the number(s) and address(es) provided on this Patient Enrollment Form for the purpose of injection reminders and any other purposes described in the preceding Patient Authorization. I confirm that I am the subscriber for the telephone number(s) provided on this Patient Enrollment Form and the authorized user for the email address(es) provided, and I agree to notify VIATRIS ADVOCATE promptly if any of my number(s) or address(es) change in the future. I understand that my wireless service provider’s message and data rates may apply.

C. MARKETING CONSENT: I would also like to receive marketing information, offers, and promotions from Viatris Inc. regarding its products, programs, and services. I agree to be contacted by email at the email address provided on this form with such information as well as with inquiries about my opinions regarding such products, programs, and services. I understand that the personal information I supply to Viatris Inc. will be shared with and among its business partners to provide me with information on Viatris-specific products, programs and services. I may cancel my participation at any time by calling 1.844.695.2667 or by following the opt-out instructions contained within the emails themselves.

USE

GLATIRAMER ACETATE INJECTION is a prescription medicine that is used to treat relapsing forms of multiple sclerosis (MS), to include clinically isolated syndrome, relapsing-remitting disease, and active secondary progressive disease, in adults. It is not known if GLATIRAMER ACETATE INJECTION is safe and effective in children under 18 years of age.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about GLATIRAMER ACETATE INJECTION?

- **Serious allergic reactions (anaphylactic reactions). Serious allergic reactions that may be life-threatening or lead to death may happen any time after you start using GLATIRAMER ACETATE INJECTION. These reactions may happen right after your first dose up to years after starting treatment with GLATIRAMER ACETATE INJECTION, even if you never had an allergic reaction before. Many reactions have happened within 1 hour of using GLATIRAMER ACETATE INJECTION. Some signs and symptoms may be the same as those of an immediate post-injection reaction. See What are the possible side effects of GLATIRAMER ACETATE INJECTION?**

Stop using GLATIRAMER ACETATE INJECTION and get emergency help right away if you have:

- **widespread rash**
- **swelling of the face, eyelids, lips, mouth, throat, or tongue**
- **sudden shortness of breath, difficulty breathing, or wheezing**
- **uncontrolled shaking (convulsions)**
- **trouble swallowing or speaking**
- **fainting, feeling dizzy or faint**

Do not take GLATIRAMER ACETATE INJECTION if you are allergic to glatiramer acetate or mannitol. Serious allergic reactions including life-threatening or anaphylactic reactions that can lead to death have happened.

Be sure to talk to your healthcare provider if you are pregnant or plan to become pregnant for them to advise if you should take glatiramer acetate during your pregnancy. Be sure to tell your healthcare provider about all of your medications.

Serious side effects may happen right after or within minutes after you inject GLATIRAMER ACETATE INJECTION at any time during your course of treatment. Call your doctor right away if you have any of these immediate post-injection reaction symptoms including redness to your cheeks or other parts of the body (flushing); chest pain; fast heartbeat; anxiety; breathing problems or tightness in your throat; or swelling, rash, hives, or itching. If you have symptoms of an immediate post-injection reaction, do not give yourself more injections until a doctor tells you to.

Chest pain may occur either as part of the immediate post-injection reaction or by itself. This type of chest pain usually lasts a few minutes or can begin around 1 month after you start using GLATIRAMER ACETATE INJECTION. Call your doctor right away if you experience chest pain.

Damage to the fatty tissue just under your skin’s surface (lipoatrophy) and, rarely, death of your skin tissue (necrosis) can happen when you use glatiramer acetate injection. Damage to the fatty tissue under your skin can cause a “dent” at the injection site that may not go away. You can reduce your chance of developing these problems by:

- following your healthcare provider’s instructions for how to use glatiramer acetate injection
- choosing a different injection area each time you use glatiramer acetate injection

Liver problems, including liver failure, can occur with GLATIRAMER ACETATE INJECTION. Call your healthcare provider right away if you have symptoms, such as nausea, loss of appetite, tiredness, dark colored urine and pale stools, yellowing of your skin or the white part of your eye, bleeding more easily than normal, confusion, or sleepiness.

Some glatiramer acetate products can be used with an optional compatible autoinjector. Compatible autoinjectors are supplied separately if available, but the availability of compatible autoinjectors may change with time. Check with your healthcare provider when you fill or refill your medicine to make sure the autoinjector you have is meant to be used with your glatiramer acetate product. If you use the wrong autoinjector, you might not get the correct dose of your medicine.

The most common side effects in studies of GLATIRAMER ACETATE INJECTION are skin problems at your injection site, including redness, pain, swelling, itching, or a lump at the site of injection, rash, shortness of breath, flushing (vasodilation), and chest pain. These are not all the possible side effects of GLATIRAMER ACETATE INJECTION. For a complete list, ask your doctor or pharmacist. Tell your doctor about any side effects that you have while taking GLATIRAMER ACETATE INJECTION.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call **1-800-FDA-1088**.

Please see accompanying Full Prescribing Information, including Boxed WARNING, Patient Information Leaflet and Instructions for Use for GLATIRAMER ACETATE INJECTION 20 mg/mL or GLATIRAMER ACETATE INJECTION 40 mg/mL.

For more information, visit glatirameracetate.com.

WhisperJECT® and the Glatiramer Acetate Injection Logo are registered trademarks of Mylan Pharmaceuticals Inc., a Viatris Company.

VIATRIS and the Viatris Logo are trademarks of Mylan Inc., a Viatris Company.

VIATRIS ADVOCATE and the Viatris Advocate Logo are registered trademarks of Mylan Inc., a Viatris Company.

© 2025 Viatris Inc. All Rights Reserved. GA-2019-0154 V8